

Sussex Research Online

Structural classification of insecticidal proteins – towards an in silico characterization of novel toxins

Article (Accepted Version)

Berry, Colin and Crickmore, Neil (2017) Structural classification of insecticidal proteins – towards an in silico characterization of novel toxins. *Journal of Invertebrate Pathology*, 142. pp. 16-22. ISSN 0022-2011

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/62164/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Accepted Manuscript

Structural classification of insecticidal proteins – towards an *in silico* characterization of novel toxins

Colin Berry, Neil Crickmore

PII: S0022-2011(16)30100-8
DOI: <http://dx.doi.org/10.1016/j.jip.2016.07.015>
Reference: YJIPA 6846

To appear in: *Journal of Invertebrate Pathology*

Received Date: 26 February 2016
Revised Date: 1 June 2016
Accepted Date: 28 July 2016



Please cite this article as: Berry, C., Crickmore, N., Structural classification of insecticidal proteins – towards an *in silico* characterization of novel toxins, *Journal of Invertebrate Pathology* (2016), doi: <http://dx.doi.org/10.1016/j.jip.2016.07.015>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Structural classification of insecticidal proteins – towards an *in silico* characterization of novel toxins.

Colin Berry^{1*} and Neil Crickmore²

¹Cardiff School of Biosciences, Cardiff University, Park Place, Cardiff CF10 3AT, UK

²School of Life Sciences, University of Sussex, Falmer, Brighton, BN1 9QG, UK

*Corresponding author: Colin Berry, Cardiff School of Biosciences, Cardiff University, Park Place, Cardiff CF10 3AT, UK, Berry@cf.ac.uk

Abstract:

The increasing rate of discovery of new toxins with potential for the control of invertebrate pests through next generation sequencing, presents challenges for the identification of the best candidates for further development. A consideration of structural similarities between the different toxins suggest that they may be functionally less diverse than their low sequence similarities might predict. This is encouraging from the perspective of being able to use computational tools to predict toxin targets from their sequences, however more structure/function data are still required to reliably inform such predictions.

Introduction:

The insecticidal toxins of *Bacillus thuringiensis*, *Lysinibacillus sphaericus*, *Photorhabdus* spp. and other bacteria represent a rich resource for the control of pest insects. The increasing rate of discovery of new toxins, driven by next-generation sequencing, will expand our arsenal of potential biocontrol agents but this, in itself, presents new challenges. Even with past rates of toxin discovery, toxins have rarely been tested against more than a few species of insects (van Frankenhuyzen, 2009) and, in the future, toxicity testing of large numbers of new toxins

against a wide range of insects will not be feasible. To facilitate the selection of toxins for study, different criteria may be applied, including identification of the toxin in a strain known (from a previous screening) to have interesting biocidal activity or relatedness to known toxins. Here we consider the prospects for a further, selective method through the prediction of activity. We highlight some of the challenges that may be encountered and propose steps that will bring us closer to this goal. Useful predictions would not only assist in the selection of toxins for development but would also have value in support of the regulatory process of biopesticide product registration, where the potential to predict off-target activities would be valuable.

The *B. thuringiensis* nomenclature system (Crickmore et al., 1998) currently contains several hundred individual sequences, divided between 74 classes of Cry toxin, 3 classes of Cyt toxin, 4 classes of Vip toxin and one SIP toxin. *L. sphaericus* strains may produce the BinA/B toxin, Mtx1, Mtx2, Mtx3, Mtx4, sphaericolysin, Cry48 and Cry49 (reviewed in (Berry, 2012)) and *Photorhabdus* strains can produce Tc toxins, PirA/B and Mcf toxins (ffrench-Constant et al., 2007). This represents a great diversity of toxins but some simplification can be achieved by considering these proteins in terms of their structural characteristics (known or predicted). Table 1 shows the toxin classes, colour-coded by sequence homology groups. As can be seen, the 3-domain Cry toxins represent the largest structural family (and also encompass the PirA/B toxin, recently shown to be equivalent to a 3-domain toxin with a dissociated domain III (Lee et al., 2015)). There is also a large group of toxins that is rich in beta-sheets with general structural similarity to aerolysin. This group includes Cry46 and toxins identified by Pfam (Bateman et al., 1999) to be members of either the Etx/Mtx2 family or the Toxin_10 family. Other groups include the Cyt toxins, the ADP-ribosyl transferase toxins Mtx1 and Vip1/2 (along with the Vip1-like Vip4 protein). Cry34 is an aerolysin like protein and with Cry35 is part of a two-component toxin (Kelker et al., 2014). Cry37, which itself is part

of a two-component toxin with Cry23, which shows structural homology with Cry34 (Rydel et al., 2001). Other toxins, which appear unrelated and have no published structures, are Cry6, Cry22, Cry55, Vip3 and Mcf. Our knowledge of the structure and function of toxins within these groups varies and it will be useful to consider the major groups separately.

The 3-domain toxins: These toxins are the best-characterised, with the first structure published in 1991 (Li et al., 1991) and with several decades of studies on the specificity and mode of action of members of this family. The steps leading to toxicity for this family are well-known and involve ingestion by the invertebrate target, solubilisation of toxin crystals in the gut, proteolytic activation by gut enzymes, one or more receptor binding step, followed by membrane insertion (Pardo-Lopez et al., 2012). Insect specificity could be mediated by any of the above steps, for example changes in proteinase activity (Loseva et al., 2002) but the most important determinants of specificity are the binding to and specificity for receptors on the surfaces of target cells.

As suggested by the name of this family, the structure of the active toxin is composed of 3 distinct structural domains. Domain I is formed from a bundle of alpha helices and is involved in pore formation by the toxin. Domain II has a beta prism structure that appears to be related to carbohydrate binding proteins and Domain III has a beta sandwich fold. Domains II and III appear to have roles in receptor binding and specificity of the toxins as demonstrated by domain swapping experiments that have altered target specificity (Lee et al., 1995; Pigott and Ellar, 2007). Bioinformatic analysis suggests that the 3 toxin domains evolve at different rates (Bravo, 1997) and this may have implications for target specificity.

Within the 3-domain toxin family, we find toxins with activity against insects in several orders, principally amongst the Lepidoptera with fewer active against the orders Diptera and Coleoptera, and with small numbers active against Hymenoptera and Hemiptera as well as toxins affecting nematodes and gastropods (reviewed recently (Palma et al., 2014a)).

Members of this family active against human cancer cells have also been reported (Ohba et al., 2009), although it is clearly unlikely that they have co-evolved with this host. However, correlation between sequence identity and target range is generally poor even when analysis is carried out at the level of the individual domains (de Maagd et al., 2001). This highlights the need for analysis at a level below that of the domains themselves. Within domain II, several exposed loops (the $\alpha 8$ loop, and loops 1, 2 and 3) have been identified as potentially important for receptor binding. The variability of these regions and their dispositions in 3 example toxins is shown in Figure 1. These surface loops are amongst the most variable in sequence and in length between individual toxins and even minor modifications have been shown to change targeting (Bravo et al., 2013). For example, Cry4Ba has no significant activity against *Culex* mosquitoes but the substitution of Asp454 in domain II loop 3 with the sequence Pro-Ala-Thr results in high toxicity to *Culex* species without reduction in toxicity towards *Aedes aegypti* (Abdullah et al., 2003). However, these loops may not be the sole domain II mediators of specificity, for example it has been shown that residues remote from these loops (illustrated in Figure 2) contribute to dipteran/lepidopteran specificity in the Cry2Aa/Cry2Ab toxins (Morse et al., 2001).

An understanding of the receptor-binding interactions of the 3-domain proteins is made complex by the diversity of putative receptor proteins for these toxins. The most commonly identified binding partners include cadherins, aminopeptidases and alkaline phosphatases (Pigott and Ellar, 2007). However, even for single toxins such as Cry1Ac and Cry4Ba, a large number of interacting proteins can be identified through proteomic studies (Bayyareddy et al., 2009; Krishnamoorthy et al., 2007) (Table 2). Although the significance of these binding interactions for toxicity is not known, we cannot discount possible physiological significance for these interactions, for example prohibitin, identified as a potential Cry4Ba binding-protein has been demonstrated to bind toxin in *Aedes* cells in culture (Kuadkitkan et al., 2012). Glycolipids are also potential receptors for 3-domain toxin binding as shown for

Cry1Ac (Garczynski and Adang, 2000), Cry2Ab (Ma et al., 2012) and may be particularly important for the nematode-active Cry5B (Barrows et al., 2007; Griffiths et al., 2005). The diversity of possible receptors and the probable involvement of more than one receptor in toxicity, adds to the challenge of predicting toxin activity.

Several studies have attempted to map regions of the toxins that may interact with receptor proteins. For example, residue Tyr445 in loop 3 of Cry1Aa domain II was identified as being important in binding to the cadherin BtR175 of *Bombyx mori* (Atsumi et al., 2005) while Val582 in domain III was shown to be important for interaction with aminopeptidase N (Atsumi et al., 2008). In parallel, there have been studies to map regions of receptors that may interact with the toxins (Pigott and Ellar, 2007) but even when a common class of receptor (eg a cadherin) is considered, there is little correspondence between regions interacting with different toxins.

The beta sheet toxins: There are several classes of toxins in Table 1 that appear to be rich in beta sheets. These include Cry37 (de Maagd et al., 2003) and the aegerolysin-like Cry34 (Kelker et al., 2014) which both form part of binary toxins (with Cry23 and Cry35 respectively, discussed below). The presence of extended beta sheets also characterises the sphaericolysin/anthrolysin family of toxins that is highly conserved across isolates from *L. sphaericus*, *B. thuringiensis*, *Bacillus cereus* and *Paenibacillus alvei* (Berry, 2012; Bourdeau et al., 2009) and appear to act as cholesterol dependent cytolysins (From et al., 2008; Nishiwaki et al., 2007). At present, the literature lacks sufficient information to allow structure/function predictions for these toxins. However, a number of proteins in Table 1 belonging to the Etx/Mtx2 family, the Toxin_10 family, along with Cry46 are rich in beta sheets and show a general fold similar to aerolysin. This group of toxins is the most numerous after the 3-domain toxins and some analysis of structure and specificity is possible. The general structure of these toxins features a head region and a tail region that features long beta

strands (available structures for these proteins are shown in Figure 3). Cry45 and Cry46 are produced by *B. thuringiensis* but have no known activity against invertebrates. However, activity against mammalian cancer cells has been demonstrated (Ohba et al., 2009). Structurally, Cry46 most closely resembles the Etx/Mtx2 family (although Pfam analysis of its primary sequence does not assign it to this family). The Etx/Mtx2 family differs from the Toxin_10 family in two clear features. In the Toxin_10 family, the head domains contain beta trefoil motifs similar to carbohydrate-binding domains and these heads are formed exclusively from the N-terminal end of the proteins. The beta trefoil domains may have a role in toxin interactions with glycoproteins or glycolipids to facilitate receptor binding or other stages of the mechanisms of action of the toxins. In contrast, in the Etx/Mtx2 family, the head lacks the beta trefoil and is composed of residues from the N-terminal region and from a further stretch of amino acids much closer to the C-terminus (before the C-terminal sequence completes the last long beta strand of the tail domain). In these features, Toxin_10 proteins resemble toxins such as the haemolytic lectin from the parasitic mushroom *Laetiporus sulphureus* (Mancheno et al., 2005), while the Etx/Mtx2 family resembles mammalian toxins such as aerolysin (Figure 3) and epsilon toxin from *Clostridium perfringens*.

When the ability of these aerolysin-like toxins to target invertebrate or mammalian cells is compared to their overall sequence, there is no obvious correlation (Figure 4A). When analysis is performed at the level of the phylogeny of individual head and tail domains (Figure 4B and C) there is still no clear correlation, indicating once again the need for deeper, subdomain analysis to be undertaken to predict activity.

Understanding the receptor binding and specificity of the Toxin_10 family is further complicated by the existence of partner proteins for many of these toxins. Cry36 is a clear exception since it is reported to act alone to kill *Diabrotica* larvae (Rupar et al., 2000). Another protein in the Toxin_10 family (41.9 kDa protein) may be encoded by *B. thuringiensis* but no toxicity has been discovered to date and no partner protein has been identified (Palma

et al., 2014b). All of the other Toxin₁₀ proteins act with their specific partner proteins to form binary toxins as follows: BinA and BinB (both Toxin₁₀ proteins)(Broadwell et al., 1990; Oei et al., 1990); Cry34 (aegerolysin-like) and Cry35 (Toxin₁₀)(Kelker et al., 2014; Masson et al., 2004); Cry48 (3-domain) and Cry49 (Jones et al., 2007). The role of each protein in these binary pairs is clearly significant to understanding the specificity of the toxins and potential binding of both components presents further challenges to prediction. The binding of both Cry48 and Cry49 to *Culex* brush border membrane fractions has been shown, with Cry49 suggested to be the principal binding component (Guo et al., 2016). In the case of the Bin toxin, BinB appears to be the major receptor-binding component in *Culex* mosquitoes but in *Anopheles* BinA also appears to be able to bind (Charles et al., 1997; Oei et al., 1992). Binding of BinB to target membranes appears to be mediated by residues at its N-terminal end (Oei et al., 1992; Romao et al., 2011; Singkhamanan et al., 2010), consistent with receptor recognition via the head domain (Srisucharitpanit et al., 2014). The BinA/BinB toxin appears to bind to a single toxin receptor, a GPI anchored α -glycosidase (Silva-Filha et al., 1999), which may simplify the investigation of binding interactions (particularly when compared to the complex receptor binding of 3-domain toxins). Both the receptor, Cqm1, from the Bin-sensitive *Culex quinquefasciatus* and the ortholog from the insensitive *Aedes aegypti* are known and a region involved in binding has been identified. This includes a Gly-Gly motif, potentially on a surface loop, which may be required for productive interaction (Ferreira et al., 2014). If binding to single receptors proves to be a general feature of this class of toxin, prediction of activity may be made more straightforward but this may also have implications for the ease with which insects may acquire resistance. A number of the Etx/Mtx2 family of Cry toxins are also reported to require a binary partner for full activity. Cry23Aa (also known as ET33) acts with Cry37 (ET34) and has activity against various coleopteran insects (Donovan et al., 2000; Ekobu et al., 2010). Cry15Aa and Cry33Aa form binary partnerships with two other proteins (40kDa and NT32KD respectively) that, due to their lack of individual activity, have not been

assigned Cry names, and share no obvious sequence similarity with any other characterized protein (Kim et al., 2003; Naimov et al., 2011).

Convergent evolution of the beta sheet toxins?:

Figure 3 indicates that three of the homology groups highlighted in Table 1 (Toxin_10, Cry46 and Etx/Mtx2) share significant sequence similarity and indeed are very similar to toxins such as aerolysin. In the aerolysin family of proteins, the conserved beta sheet dominated “aerolysin fold” is believed to adopt a barrel conformation within a membrane with the associated domains primarily having a binding role (Szczesny et al., 2011). It is conceivable that despite the diversity in primary sequence, many of the non 3-domain Cry toxins could share significant structural/functional homology. This would take the form of the aerolysin tail fold associated with a head domain involved in targeting the protein to a particular receptor. This head domain could be part of the same protein as the aerolysin fold or come from an associated binary partner. A number of Cry and Cyt toxins have acquired a ricin-like beta-trefoil carbohydrate-binding domain and while it is tempting to speculate that this could have given these proteins novel binding activities, there is currently no evidence that this is the case (reviewed by (Adang et al., 2014)).

The future for *in silico* analyses:

The ultimate objective of such work is to be able to predict the likely specificity of a toxin from primary sequence data. While the computational power to be able to achieve such a goal is available, the underlying data required to inform such analyses are still lacking. Significant progress has been made in recent years in the elucidation of new toxin structures and the potential to derive reliable structures from modelling approaches is discussed elsewhere in this issue (Berry and Board, 2016). Similarly, while there are a lot of data available on the target specificity of individual toxins (Palma et al., 2014a; van Frankenhuyzen, 2009), many of

these are contradictory, thus compounding efforts to derive meaningful associations. The area where least information is known concerns the interaction of the toxin with the target cell (Vachon et al., 2012). As discussed above, even when putative receptors are identified, determining which interactions are crucial for toxicity is far from straightforward. There is a need to elucidate more structures for confirmed receptors, and ideally toxin-receptor complexes, which can then lead to *in silico* predictions of toxin-receptor interactions. A number of studies have used docking analysis to indicate the likely interaction between a Cry toxin and its putative receptor (Ahmad et al., 2015; Tajne et al., 2012; Zhao et al., 2012). Without a detailed understanding of which interactions are crucial for toxicity, such studies are likely to throw up many false positives, indeed one report predicts that Cry1Ac could have activity against cattle (Ebenezer et al., 2013), a prediction that is not supported by experimental observations.

In summary, whilst our understanding of toxin structure is rapidly progressing, and might suggest that the large family of toxins is less diverse than was thought, we are still a long way from the goal of being able to match toxins and hosts based on primary sequence data generated from genome sequencing.

Table1: Toxins and their homology groups

Toxins of *B. thuringiensis*, *L. sphaericus* and *Photorhabdus* spp. are shown with colouring to indicate homology groups: light blue = 3-domain toxins; peach = Etx/Mtx2 toxins; pink = Toxin_10 family proteins; violet = Cyt toxins; khaki = aegerolysin toxins; grey = ADP ribosyl transferase-related proteins; toxins not falling in to these groups are coloured differently.

Cry1	Cry21	Cry41	Cry61	Vip3
Cry2	Cry22	Cry42	Cry62	Vip4
Cry3	Cry23	Cry43	Cry63	BinA
Cry4	Cry24	Cry44	Cry64	BinB
Cry5	Cry25	Cry45	Cry65	Mtx1
Cry6	Cry26	Cry46	Cry66	Mtx2
Cry7	Cry27	Cry47	Cry67	Mtx3
Cry8	Cry28	Cry48	Cry68	Mtx4
Cry9	Cry29	Cry49	Cry69	Sphaericolysin
Cry10	Cry30	Cry50	Cry70	PirA
Cry11	Cry31	Cry51	Cry71	PirB
Cry12	Cry32	Cry52	Cry72	Mcf
Cry13	Cry33	Cry53	Cry73	
Cry14	Cry34	Cry54	Cry74	
Cry15	Cry35	Cry55	Cyt1	
Cry16	Cry36	Cry56	Cyt2	
Cry17	Cry37	Cry57	Cyt3	
Cry18	Cry38	Cry58	Sip	
Cry19	Cry39	Cry59	Vip1	
Cry20	Cry40	Cry60	Vip2	

Table 2: Potential receptors for 3-domain toxins.

Proteins discovered through proteomics as potential binding partners for two 3-domain toxins, Cry1Ac and Cry4Ba are shown.

Cry4Ba (Bayyareddy et al., 2009)	Cry1Ac (Krishnamoorthy et al., 2007)
Cadherin	Cadherin
Alkaline phosphatases (3)	Alkaline phosphatases
ATPase	ATPases
Actin	Actin
Serine and metallo peptidases	Aminopeptidases
Prohibitin	Desmocollin-like protein
Mitoporin	
Flotillin-1	
ATP synthase	

Figure Legends

Figure 1: Domain II loops

The structures of Cry1Aa, Cry2Aa and Cry3Aa (PDB codes 1CIY, 1I5P and 1DLC respectively) are shown with domains I and III shown as yellow and brown ribbons while domain II is shown as a surface representation with the exposed regions of the α 8 loop, and loops 1, 2 and 3 coloured cyan, orange, green and magenta, respectively.

Figure 2: Residues implicated in Cry2A specificity

The Cry2Aa structure (PDB 1I5P) is shown with domain II in grey. Residues shown to contribute to lepidopteran specificity are coloured blue while those involved in dipteran specificity are shown in magenta.

Figure 3: Structures of beta sheet toxins

The structures of the insecticidal toxins BinB (PDB 3WA1 (Srisucharitpanit et al., 2014)), Cry35 (PDB 4JP0 (Kelker et al., 2014)), Cry23 (PDB 4RHZ), Cry45 (PDB 2D42 (Akiba et al., 2006)), Cry46 (PDB 2ZTB (Akiba et al., 2009)) and Cry51 (PDB 4PKM (Xu et al., 2015)) are shown along with the structures of the haemolytic toxin from *L. sulphureus* (Lsulph: PDB 1W3A (Mancheno et al., 2005) and proaerolysin (PDB 1PRE (Parker et al., 1994)). Head regions are coloured cyan, tails blue and the extra domain in proaerolysin in green.

Figure 4: Phylogenetic relationship of insecticidal and mammalian-active beta sheet toxins

The amino acid sequences of the toxins were compared using MEGA6 and phylogenetic trees were built using the maximum likelihood algorithm. For those toxins where the head and tail domains are discontinuous, the separate regions encoding each domain were combined and analysed as a contiguous sequence. Those toxins with known activity against mammals are highlighted in bold red.

References:

Abdullah, M. A., Alzate, O., Mohammad, M., McNall, R. J., Adang, M. J., Dean, D. H., 2003.

Introduction of *Culex* toxicity into *Bacillus thuringiensis* Cry4Ba by protein engineering.

Appl Environ Microbiol. 69, 5343-5353.

Adang, M. J., Crickmore, N., Jurat-Fuentes, J. L. Eds.), 2014. Diversity of *Bacillus thuringiensis* Crystal Toxins and Mechanism of Action. Elsevier.

- Ahmad, A., Javed, M. R., Rao, A. Q., Khan, M. A., Ahad, A., Din, S. U., Shahid, A. A., Husnain, T., 2015. *In-Silico* Determination of Insecticidal Potential of Vip3Aa-Cry1Ac Fusion Protein Against Lepidopteran Targets Using Molecular Docking. *Frontiers in plant science*. 6, 1081.
- Akiba, T., Abe, Y., Kitada, S., Kusaka, Y., Ito, A., Ichimatsu, T., Katayama, H., Akao, T., Higuchi, K., Mizuki, E., Ohba, M., Kanai, R., Harata, K., 2009. Crystal structure of the parasporin-2 *Bacillus thuringiensis* toxin that recognizes cancer cells. *J Mol Biol*. 386, 121-133.
- Akiba, T., Higuchi, K., Mizuki, E., Ekino, K., Shin, T., Ohba, M., Kanai, R., Harata, K., 2006. Nontoxic crystal protein from *Bacillus thuringiensis* demonstrates a remarkable structural similarity to beta-pore-forming toxins. *Proteins*. 63, 243-248.
- Atsumi, S., Inoue, Y., Ishizaka, T., Mizuno, E., Yoshizawa, Y., Kitami, M., Sato, R., 2008. Location of the *Bombyx mori* 175kDa cadherin-like protein-binding site on *Bacillus thuringiensis* Cry1Aa toxin. *FEBS J*. 275, 4913-4926.
- Atsumi, S., Mizuno, E., Hara, H., Nakanishi, K., Kitami, M., Miura, N., Tabunoki, H., Watanabe, A., Sato, R., 2005. Location of the *Bombyx mori* aminopeptidase N type 1 binding site on *Bacillus thuringiensis* Cry1Aa toxin. *Appl Environ Microbiol*. 71, 3966-3977.
- Barrows, B. D., Griffiths, J. S., Aroian, R. V., 2007. Resistance is non-futile: resistance to Cry5B in the nematode *Caenorhabditis elegans*. *J Invertebr Pathol*. 95, 198-200.
- Bateman, A., Birney, E., Durbin, R., Eddy, S. R., Finn, R. D., Sonnhammer, E. L., 1999. Pfam 3.1: 1313 multiple alignments and profile HMMs match the majority of proteins. *Nucleic Acids Res*. 27, 260-262.
- Bayyareddy, K., Andacht, T. M., Abdullah, M. A., Adang, M. J., 2009. Proteomic identification of *Bacillus thuringiensis* subsp. *israelensis* toxin Cry4Ba binding proteins in midgut membranes from *Aedes* (*Stegomyia*) *aegypti* Linnaeus (Diptera, Culicidae) larvae. *Insect Biochem Mol Biol*. 39, 279-286.

- Berry, C., 2012. The bacterium, *Lysinibacillus sphaericus*, as an insect pathogen. J. Invertebr. Pathol. 109, 1-10.
- Berry, C., Board, J., 2016. Insights into the structures of non-3-domain toxins through structural modelling. J. Invertebr. Pathol. JIP-16-21.
- Bourdeau, R. W., Malito, E., Chenal, A., Bishop, B. L., Musch, M. W., Villereal, M. L., Chang, E. B., Mosser, E. M., Rest, R. F., Tang, W. J., 2009. Cellular functions and X-ray structure of anthrolysin O, a cholesterol-dependent cytolysin secreted by *Bacillus anthracis*. J Biol Chem. 284, 14645-14656.
- Bravo, A., 1997. Phylogenetic relationships of *Bacillus thuringiensis* delta-endotoxin family proteins and their functional domains. J. Bacteriol. 179, 2793-2801.
- Bravo, A., Gomez, I., Porta, H., Garcia-Gomez, B. I., Rodriguez-Almazan, C., Pardo, L., Soberon, M., 2013. Evolution of *Bacillus thuringiensis* Cry toxins insecticidal activity. Microb Biotechnol. 6, 17-26.
- Broadwell, A. H., Baumann, L., Baumann, P., 1990. Larvicidal properties of the 42 and 51 kilodalton *Bacillus sphaericus* proteins expressed in different bacterial hosts: evidence for a binary toxin. Curr. Microbiol. 21, 361-366.
- Charles, J.-F., Silva-Filha, M. H., Nielsen-LeRoux, C., Humphreys, M. J., Berry, C., 1997. Binding of the 51- and 42-kDa individual components from the *Bacillus sphaericus* crystal toxin to mosquito larval midgut membranes from *Culex* and *Anopheles* sp. (Diptera: Culicidae). FEMS Microbiol. Lett. 156, 153-159.
- Crickmore, N., Zeigler, D. R., Feitelson, J., Schnepf, E., Van Rie, J., Lereclus, D., Baum, J., Dean, D. H., 1998. Revision of the Nomenclature for the *Bacillus thuringiensis* Pesticidal Crystal Proteins. Microbiol. Mol. Biol. Rev. 62, 807-813.
- de Maagd, R. A., Bravo, A., Berry, C., Crickmore, N., Schnepf, H. E., 2003. Structure, diversity, and evolution of protein toxins from spore-forming entomopathogenic bacteria. Annu Rev Genet. 37, 409-433.

- de Maagd, R. A., Bravo, A., Crickmore, N., 2001. How *Bacillus thuringiensis* has evolved specific toxins to colonize the insect world. *Trends Genet.* 17, 193-9.
- Donovan, W. P., Donovan, J. C., Slaney, A. C., *Bacillus thuringiensis* cryET33 and cryET34 compositions and uses therefor. Vol. . Monsanto Company, 2000.
- Ebenezer, K. S., Nachimuthu, R., Thiagarajan, P., Velu, R. K., 2013. in silico identification of cross affinity towards Cry1Ac pesticidal protein with receptor enzyme in *Bos taurus* and sequence, structure analysis of crystal proteins for stability. *Bioinformation.* 9, 792-795.
- Ekobu, M., Solera, M., Kyamanywa, S., Mwanga, R. O., Odongo, B., Ghislain, M., Moar, W. J., 2010. Toxicity of seven *Bacillus thuringiensis* Cry proteins against *Cylas puncticollis* and *Cylas brunneus* (Coleoptera: Brentidae) using a novel artificial diet. *Journal of economic entomology.* 103, 1493-1502.
- Ferreira, L. M., Romao, T. P., Nascimento, N. A., Costa, M. D., Rezende, A. M., de-Melo-Neto, O. P., Silva-Filha, M. H., 2014. Non conserved residues between Cqm1 and Aam1 mosquito alpha-glucosidases are critical for the capacity of Cqm1 to bind the Binary (Bin) toxin from *Lysinibacillus sphaericus*. *Insect Biochem Mol Biol.*
- ffrench-Constant, R. H., Dowling, A., Waterfield, N. R., 2007. Insecticidal toxins from *Photorhabdus* bacteria and their potential use in agriculture. *Toxicon.* 49, 436-451.
- From, C., Granum, P. E., Hardy, S. P., 2008. Demonstration of a cholesterol-dependent cytolysin in a noninsecticidal *Bacillus sphaericus* strain and evidence for widespread distribution of the toxin within the species. *FEMS Microbiol Lett.* 286, 85-92.
- Garczynski, S., Adang, M. J., Investigations of *Bacillus thuringiensis* Cry1 toxin receptor structure and function. In: J.-F. Charles, et al., Eds.), *Entomopathogenic Bacteria: From Laboratory to Field Application.* Kluwer Academic Publishers, Dordrecht, The Netherlands, 2000, pp. 181-197.

- Griffitts, J. S., Haslam, S. M., Yang, T., Garczynski, S. F., Mulloy, B., Morris, H., Cremer, P. S., Dell, A., Adang, M. J., Aroian, R. V., 2005. Glycolipids as receptors for *Bacillus thuringiensis* crystal toxin. *Science*. 307, 922-925.
- Guo, Q. Y., Hu, X. M., Cai, Q. X., Yan, J. P., Yuan, Z. M., 2016. Interaction of *Lysinibacillus sphaericus* Cry48Aa/Cry49Aa toxin with midgut brush-border membrane fractions from *Culex quinquefasciatus* larvae. *Insect Mol Biol*.
- Jones, G. W., Nielsen-Leroux, C., Yang, Y., Yuan, Z., Dumas, V. F., Monnerat, R. G., Berry, C., 2007. A new Cry toxin with a unique two-component dependency from *Bacillus sphaericus*. *FASEB J*. 21, 4112-4120.
- Kelker, M. S., Berry, C., Evans, S. L., Pai, R., McCaskill, D. G., Wang, N. X., Russell, J. C., Baker, M. D., Yang, C., Pflugrath, J. W., Wade, M., Wess, T. J., Narva, K. E., 2014. Structural and Biophysical Characterization of *Bacillus thuringiensis* Insecticidal Proteins Cry34Ab1 and Cry35Ab1. *PLoS One*. 9, e112555.
- Kim, H. S., Saitoh, H., Yamashita, S., Akao, T., Park, Y. S., Maeda, M., Tanaka, R., Mizuki, E., Ohba, M., 2003. Cloning and characterization of two novel crystal protein genes from a *Bacillus thuringiensis* serovar *dakota* strain. *Current microbiology*. 46, 33-8.
- Krishnamoorthy, M., Jurat-Fuentes, J. L., McNall, R. J., Andacht, T., Adang, M. J., 2007. Identification of novel CryIaC binding proteins in midgut membranes from *Heliothis virescens* using proteomic analyses. *Insect Biochemistry and Molecular Biology*. 37, 189-201.
- Kuadkitkan, A., Smith, D. R., Berry, C., 2012. Investigation of the Cry4B-prohibitin interaction in *Aedes aegypti* cells. *Curr Microbiol*. 65, 446-454.
- Lee, C. T., Chen, I. T., Yang, Y. T., Ko, T. P., Huang, Y. T., Huang, J. Y., Huang, M. F., Lin, S. J., Chen, C. Y., Lin, S. S., Lightner, D. V., Wang, H. C., Wang, A. H., Wang, H. C., Hor, L. I., Lo, C. F., 2015. The opportunistic marine pathogen *Vibrio parahaemolyticus* becomes virulent by acquiring a plasmid that expresses a deadly toxin. *Proc Natl Acad Sci U S A*.

- Lee, M. K., Young, B. A., Dean, D. H., 1995. Domain III exchanges of *Bacillus thuringiensis* CryIA toxins affect binding to different gypsy moth midgut receptors. *Biochem Biophys Res Commun.* 216, 306-312.
- Li, J. D., Carroll, J., Ellar, D. J., 1991. Crystal structure of insecticidal delta-endotoxin from *Bacillus thuringiensis* at 2.5 Å resolution. *Nature.* 353, 815-821.
- Loseva, O., Ibrahim, M., Candas, M., Koller, C. N., Bauer, L. S., Bulla, L. A., Jr., 2002. Changes in protease activity and Cry3Aa toxin binding in the Colorado potato beetle: implications for insect resistance to *Bacillus thuringiensis* toxins. *Insect Biochem Mol Biol.* 32, 567-77.
- Ma, G., Rahman, M. M., Grant, W., Schmidt, O., Asgari, S., 2012. Insect tolerance to the crystal toxins Cry1Ac and Cry2Ab is mediated by the binding of monomeric toxin to lipophorin glycolipids causing oligomerization and sequestration reactions. *Developmental and Comparative Immunology.* 37, 184-192.
- Mancheno, J. M., Tateno, H., Goldstein, I. J., Martinez-Ripoll, M., Hermoso, J. A., 2005. Structural analysis of the *Laetiporus sulphureus* hemolytic pore-forming lectin in complex with sugars. *J Biol Chem.* 280, 17251-17259.
- Masson, L., Schwab, G., Mazza, A., Brousseau, R., Potvin, L., Schwartz, J. L., 2004. A novel *Bacillus thuringiensis* (PS149B1) containing a Cry34Ab1/Cry35Ab1 binary toxin specific for the western corn rootworm *Diabrotica virgifera virgifera* LeConte forms ion channels in lipid membranes. *Biochemistry.* 43, 12349-12357.
- Morse, R. J., Yamamoto, T., Stroud, R. M., 2001. Structure of Cry2Aa suggests an unexpected receptor binding epitope. *Structure.* 9, 409-417.
- Naimov, S., Valkova, R., Dukiandjiev, S., Minkov, I., de Maagd, R. A., 2011. Carboxy-terminal extension effects on crystal formation and insecticidal properties of Cry15Aa. *J Invertebr Pathol.* 108, 56-58.

- Nishiwaki, H., Nakashima, K., Ishida, C., Kawamura, T., Matsuda, K., 2007. Cloning, Functional Characterization, and Mode of Action of a Novel Insecticidal Pore-Forming Toxin, Sphaericolysin, Produced by *Bacillus sphaericus*. *Appl. Environ. Microbiol.* 73, 3404-3411.
- Oei, C., Hindley, J., Berry, C., 1990. An analysis of the genes encoding the 51.4- and 41.9-kDa toxins of *Bacillus sphaericus* 2297 by deletion mutagenesis: the construction of fusion proteins. *FEMS Microbiol. Lett.* 72, 265-274.
- Oei, C., Hindley, J., Berry, C., 1992. Binding of purified *Bacillus sphaericus* binary toxin and its deletion derivatives to *Culex quinquefasciatus* gut: elucidation of functional binding domains. *J. Gen. Microbiol.* 138, 1515-1526.
- Ohba, M., Mizuki, E., Uemori, A., 2009. Parasporin, a new anticancer protein group from *Bacillus thuringiensis*. *Anticancer Res.* 29, 427-433.
- Palma, L., Muñoz, D., Berry, C., Murillo, J., Caballero, P., 2014a. *Bacillus thuringiensis* toxins: an overview of their biocidal activity. *Toxins.* 6, 3296-3325.
- Palma, L., Muñoz, D., Berry, C., Murillo, J., Caballero, P., 2014b. Draft genome sequences of two *Bacillus thuringiensis* strains and characterization of a putative 41.9-kDa insecticidal toxin. *Toxins.* 6, 1490-1504.
- Pardo-Lopez, L., Soberon, M., Bravo, A., 2012. *Bacillus thuringiensis* insecticidal 3-domain Cry toxins: Mode of action, insect resistance and consequences for crop protection. *FEMS Microbiol Rev.*
- Parker, M. W., Buckley, J. T., Postma, J. P., Tucker, A. D., Leonard, K., Pattus, F., Tsernoglou, D., 1994. Structure of the *Aeromonas* toxin proaerolysin in its water-soluble and membrane-channel states. *Nature.* 367, 292-295.
- Pigott, C. R., Ellar, D. J., 2007. Role of receptors in *Bacillus thuringiensis* crystal toxin activity. *Microbiol Mol Biol Rev.* 71, 255-271.

- Romao, T. P., de-Melo-Neto, O. P., Silva-Filha, M. H., 2011. The N-terminal third of the BinB subunit from *Bacillus sphaericus* binary toxin is sufficient for its interaction with midgut receptors in *Culex quinquefasciatus*. FEMS microbiology letters.
- Rupar, M. J., Donovan, W. P., Chu, C.-R., Pease, E., Slaney, A. C., Malvar, T. M., Baum, J. A., Coleopteran-toxic polypeptide compositions and insect-resistant transgenic plants. Vol. , 2000.
- Rydel, T., Sharamitaro, J., Brown, G. R., Gouzov, V., Searle, J., Sturman, E., Thoma, R., Grays, K., Isaac, B., The crystal structure of a coleopteran insect-active binary B.t. protein toxin complex at 2.5 Å resolution. Annual meeting of the American Crystallographic Association, Los Angeles, 2001.
- Silva-Filha, M. H., Nielsen-LeRoux, C., Charles, J.-F., 1999. Identification of the receptor for *Bacillus sphaericus* crystal toxin in the brush border membrane of the mosquito *Culex pipiens* (Diptera: Culicidae). Insect Biochem. Mol. Biol. 29, 711-721.
- Singkhamanan, K., Promdonkoy, B., Chaisri, U., Boonserm, P., 2010. Identification of amino acids required for receptor binding and toxicity of the *Bacillus sphaericus* binary toxin. FEMS Microbiol Lett. 303, 84-91.
- Srisucharitpanit, K., Yao, M., Promdonkoy, B., Chimnaronk, S., Tanaka, I., Boonserm, P., 2014. Crystal structure of BinB: A receptor binding component of the binary toxin from *Lysinibacillus sphaericus*. Proteins. 82, 2703-2712.
- Szczesny, P., Iacovache, I., Muszewska, A., Ginalska, K., van der Goot, F. G., Grynberg, M., 2011. Extending the aerolysin family: from bacteria to vertebrates. PloS one. 6, e20349.
- Tajne, S., Sanam, R., Gundla, R., Gandhi, N. S., Mancera, R. L., Boddupally, D., Vudem, D. R., Khareedu, V. R., 2012. Molecular modeling of Bt Cry1Ac (DI-DII)-ASAL (*Allium sativum* lectin)-fusion protein and its interaction with aminopeptidase N (APN) receptor of *Manduca sexta*. Journal of molecular graphics & modelling. 33, 61-76.

- Vachon, V., Laprade, R., Schwartz, J. L., 2012. Current models of the mode of action of *Bacillus thuringiensis* insecticidal crystal proteins: A critical review. J Invertebr Pathol. 111, 1-12.
- van Frankenhuyzen, K., 2009. Insecticidal activity of *Bacillus thuringiensis* crystal proteins. J Invertebr Pathol. 101, 1-16.
- Xu, C., Chinte, U., Chen, L., Yao, Q., Meng, Y., Zhou, D., Bi, L. J., Rose, J., Adang, M. J., Wang, B. C., Yu, Z., Sun, M., 2015. Crystal structure of Cry51Aa1: A potential novel insecticidal aerolysin-type beta-pore-forming toxin from *Bacillus thuringiensis*. Biochem Biophys Res Commun. 462, 184-189.
- Zhao, X. M., Zhou, P. D., Xia, L. Q., 2012. Homology modeling of mosquitocidal Cry30Ca2 of *Bacillus thuringiensis* and its molecular docking with N-acetylgalactosamine. Biomedical and environmental sciences : BES. 25, 590-6.

Figure 1




Figure2

Figure 2




Figure3

Figure 3






Figure4

Figure 4


A Overall



B Head




C Tail



Sequence diversity

Cry1
PirA
PirB
Cyt1
Cry6
Cry22
Cry34
Cry35
Cry46
Cry51
Vip1
Vip2
Vip3
Mcf
Sphaericolysin

Structural diversity



Highlights

An overview of the structural classes of invertebrate active toxins is presented along with discussion of their structure-function relationships. The prospects for predictive tools to select toxins against specific targets, based on their sequences is discussed and deficits in our present knowledge are highlighted, indicating several important areas for new research